

SPECTROSCOPIC CONFIRMATION OF MULTIPLE RED GALAXY-GALAXY MERGERS IN MS 1054-03 ($Z = 0.83$)^{1,2,3}

KIM-VY H. TRAN⁴, PIETER VAN DOKKUM⁵, MARIJN FRANX⁶, GARTH D. ILLINGWORTH⁷, DANIEL
 D. KELSON⁸, & NATASCHA M. FÖRSTER SCHREIBER⁹
Draft version February 2, 2008

ABSTRACT

We present follow-up spectroscopy of the galaxy cluster MS 1054-03 ($z = 0.83$) confirming that at least six of the nine merging galaxy pairs identified by van Dokkum et al. (1999) are indeed bound systems: they have projected separations of $R_s < 10h^{-1}$ kpc and relative line-of sight velocities of $\Delta v < 165$ km s⁻¹. For the remaining three pairs, we were unable to obtain redshifts of both constituent galaxies. To identify a more objective sample of merging systems, we select bound red galaxy pairs ($R_s \leq 30h^{-1}$ kpc, $\Delta v \leq 300$ km s⁻¹) from our sample of 121 confirmed cluster members: galaxies in bound red pairs make up $15.7 \pm 3.6\%$ of the cluster population. The $(B - K_s)$ color-magnitude diagram shows that the pair galaxies are as red as the E/S0 members and have a homogeneous stellar population. The red pair galaxies span a large range in luminosity and internal velocity dispersion to include some of the brightest, most massive members ($L > L^*$, $\sigma_{1D} > 200$ km s⁻¹); these bound galaxy pairs must evolve into E/S0 members by $z \sim 0.7$. These results combined with MS1054's high merger fraction and reservoir of likely future mergers indicates that most, if not all, of its early-type members evolved from (passive) galaxy-galaxy mergers at $z \lesssim 1$.

Subject headings: galaxies: clusters: individual (MS 1054-03) – galaxies: elliptical and lenticular, cD – galaxies: fundamental parameters – galaxies: evolution

1. INTRODUCTION

Whether galaxy-galaxy merging plays a prominent role in the formation of early-type galaxies in clusters remains an open question. The expected merger rate in rich, virialized clusters is low because the relative velocities of members are too high (*e.g.* Makino & Hut 1997). Observational studies are consistent with this idea as the fraction of galaxies in merging systems is low in clusters at $z \lesssim 0.3$ (*e.g.* Dressler 1980; Couch et al. 1998). In this context, the discovery by van Dokkum et al. (1999, hereafter vD99) that 17% of the galaxies in the rich cluster MS 1054-03 ($z = 0.83$) are in merging systems was a surprise. vD99 argue that because MS1054 has not yet virialized, the high merger rate in this cluster is “permitted.”

The main uncertainty in the vD99 study is that the merging systems were identified by eye, and identifying bound galaxy pairs in a more objective manner is needed to isolate a representative sample of merging systems. Such a sample provides better constraints on the end-product of the mergers and how galaxy-galaxy merging changes the global cluster population. For example, significant amounts of merging between red members will alter the cluster's stellar mass function. The most robust approach for selecting these “future” mergers is via

a combination of projected galaxy separation and relative line-of-sight velocity, but it does require a great deal of spectroscopy.

Motivated by these issues, we revisit the issue of galaxy-galaxy merging in MS1054 using an expanded sample of spectroscopically confirmed cluster members. In this paper, we use $(\Omega_M, \Omega_\Lambda, H_0)$ of (0.3, 0.7, 70); at $z = 0.83$, this gives a physical scale of 7.6 kpc per arcsecond.

2. SUMMARY OF OBSERVATIONS

2.1. Optical and Near-Infrared Imaging

MS1054 was part of a large program carried out by our group to study galaxy clusters at intermediate redshift. As part of our observing campaign, we obtained wide-field ($R_{BCG} \sim 1h^{-1}$ Mpc), high resolution imaging of MS1054 with HST/WFPC2 in F606W and F814W. Details of the image reduction and morphological classification are in van Dokkum et al. (2000a).

In addition, deep wide-field ground-based imaging of MS1054 in UBV and J_sHK_s was taken on the VLT with FORS1 and ISAAC, respectively. Details of the image reduction and photometry can be found in Förster Schreiber et al. (2005). To summarize, all of the images were convolved to a common spatial resolution of

¹ Based on observations with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

² Based on observations obtained at the W. M. Keck Observatory, which is operated jointly by Caltech and the University of California.

³ Based on observations obtained at the the European Southern Observatory, Paranal, Chile.

⁴ Institute for Astronomy, ETH Zürich, CH-8093 Zürich, Switzerland, vy@phys.ethz.ch

⁵ Department of Astronomy, Yale University, New Haven, CT 06520-8101

⁶ Leiden Observatory, P.O. Box 9513, 2300 RA Leiden, The Netherlands

⁷ University of California Observatories/Lick Observatory, University of California, Santa Cruz, CA 95064

⁸ Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA, 91101

⁹ Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstrasse, 85748 Garching bei München, Germany

FWHM = 0.69". Objects then were detected using SExtractor v2.2.2 (Bertin & Arnouts 1996) and colors measured using custom isophotal apertures.

2.2. Keck/LRIS Spectroscopy

To determine cluster membership, we observed MS1054 using LRIS (Oke et al. 1995) on the Keck Telescope from February 1997 to June 2001. During 6 observing runs, 21 multi-slit masks were used to measure redshifts and internal velocity dispersions for I -selected targets; one mask (June 2001) specifically targeted classified mergers. Details on how the spectra were reduced, redshifts determined, and spectral indices measured ([OII] λ 3727, H δ , & H γ) are in Tran et al. (1999, 2003). The typical error associated with the redshifts is $\delta v \sim 30 \text{ km s}^{-1}$.

Considering only galaxies that fall on the HST/WFPC2 mosaic, we have 121 confirmed cluster members that span a range in luminosity ($-22.9 < M_{Be} - 5 \log h < -18.7$); for these galaxies, we have measured structural parameters, colors, and morphologies. Our analysis also includes measured internal velocity dispersions (σ_{1D}) for 26 members (Kelson et al. 2001; Wuyts et al. 2004).

3. VISUALLY IDENTIFIED GALAXY MERGERS

We have obtained follow-up spectroscopy for the original 13 mergers reported by vD99 to determine if they are truly bound systems. To clarify, WFPC2 imaging showed that these mergers were made of 9 close pairs and 4 (single) morphologically irregular galaxies; we refer to Postman et al. (2005, Fig. 8) for higher resolution imaging from the Advanced Camera for Surveys (ACS) of the same systems. We focus here on the 9 close pairs.

Redshifts of the constituent galaxies for 6 of the 9 close pairs confirms that these are bound systems: they have projected separations of $R_s < 10h^{-1} \text{ kpc}$ and relative velocities of $\Delta v < 165 \text{ km s}^{-1}$. To confirm that their relative velocities are not due to chance projections between members, we compare their Δv to that of a random (normalized) cluster sample where the latter is generated by drawing 10^4 galaxy pairs from the full redshift sample of cluster members (Fig. 1, left). A Kolmogorov-Smirnov test shows the two distributions differ at the $> 99.7\%$ level.

For the remaining 3 close pairs, we were unable to obtain redshifts for the companion galaxies: one companion was not targeted, and the spectra of the other 2 were too faint to measure a redshift. However, we stress that all 6 of the close pairs for which we have measured relative velocities confirm that they are merging systems. Although we were unable to obtain relative velocities for the remaining 3 close pairs from vD99, it is very likely that they too are bound systems.

4. OBJECTIVELY IDENTIFIED GALAXY PAIRS

4.1. Counting Close Galaxy Pairs

As a quantitative measure of the excess of close pairs in MS1054, we follow van Dokkum et al. (2000b) and examine the distribution of galaxy-galaxy separations in the WFPC2 photometric catalog. For each galaxy, we count the number of objects N that lie at a given distance R_s and then weight by the surface area to obtain the surface density. By summing over all galaxies for a given galaxy-galaxy separation, we then have the relative galaxy density

δ as a function of R_s . For simplicity, we bin the sample into rings defined by R_{max} and R_{min} where R_s is the average of these two values. Thus for a given ring, we have:

$$\delta = \frac{\sum_i^j (N^i)}{\pi(R_{max}^2 - R_{min}^2)} \quad (1)$$

where j is the total number of galaxies. If the galaxies are distributed uniformly, the distribution of δ as a function of R_s will be flat.

Because we are primarily interested in the red cluster members, we consider only the 142 galaxies from the WFPC2 catalog with $(R_{606} - I_{814}) > 1.7$ and $I_{814} < 23$ mags; for reference, the brightest cluster galaxy (BCG) has $(R_{606} - I_{814}) = 2.13$ and $I_{814} = 19.5$ mags. Our color and magnitude selection are effective at isolating red cluster members: we have measured redshifts for 91 of these red galaxies and 89 (98%) are confirmed cluster members.

Figure 1 (middle) shows the relative galaxy density δ as a function of increasing R_s for these red galaxies. Since the high galaxy density in the cluster core can increase δ for small separations, we exclude the inner $R_{BCG} < 200h^{-1} \text{ kpc}$ region. The δ distribution is essentially flat except for a strong peak at $R_s < 10h^{-1} \text{ kpc}$.

To further investigate the issue of projected versus bound pairs, we consider the 5 red pairs with $R_s < 10h^{-1} \text{ kpc}$ and 7 red pairs with $10 \leq R_s < 20h^{-1} \text{ kpc}$ at $R_{BCG} > 200h^{-1} \text{ kpc}$. From the average relative density of pairs with $R_s > 50h^{-1} \text{ kpc}$, we estimate that 4/5 and 3/7 are bound pairs with the remainder being chance projections. We test this result with our spectroscopy. We find that all 5 red galaxy pairs with $R_s < 10h^{-1} \text{ kpc}$ are bound; note that all 5 also were identified by vD99 as mergers. Spectroscopy for 5 of the 7 red pairs with $10 \leq R_s < 20h^{-1} \text{ kpc}$ shows that only 1 pair is bound.

4.2. Defining a Sample of Bound Red Galaxy Pairs

As demonstrated above, spectroscopy is critical for determining whether photometric pairs ($R_s > 10h^{-1} \text{ kpc}$) are bound systems. With this in mind, we now define a spectroscopically confirmed sample of bound red galaxy pairs. First we take the 142 red galaxies and make each one into a pair with every other red galaxy. We then consider only the red pairs that are made of confirmed cluster members and measure their relative velocities and projected separations (Fig. 1, right).

Guided by studies on merging pairs (Patton et al. 2000; Lin et al. 2004), we select bound pairs as members with projected separations of $R_s \leq 30h^{-1} \text{ kpc}$ and relative velocities of $\Delta v \leq 300 \text{ km s}^{-1}$: there are 10 such systems (Fig. 2). Thus the fraction of red merging galaxies in MS1054 is $15.7 \pm 3.6\%$ (19/121). With this definition, a single galaxy can contribute to two separate bound pairs; there is one such triplet system in our pair sample. Note that this differs from vD99 where merging pairs were defined to be a single system. We also verified that our spectroscopic sample is not biased towards red galaxies with small projected separations: to $I_{814} = 22.5$ mags, the spectroscopic sample is $79 \pm 5\%$ complete compared to $69 \pm 14\%$ complete for red galaxies with a neighbor within $R_s \leq 30h^{-1} \text{ kpc}$.

As with all observational studies on galaxy pairs, we cannot say with absolute certainty that our 10 bound red pairs will merge into a single system by $z \sim 0$. However, detailed simulations of major mergers find that the constituent galaxies merge in $0.5 - 2$ Gyr (e.g. Barnes 1988; Mihos & Hernquist 1994; Springel 2000; Jonsson et al. 2005). In addition, Patton et al. (2000) showed that the typical merging timescale for galaxy pairs with $R_s \leq 30$ kpc and $\Delta v \leq 500$ km s $^{-1}$ is 0.5 Gyr: given our bound pairs have even smaller projected separations and relative velocities ($R_s \leq 25h^{-1}$ kpc, $\Delta v \leq 300$ km s $^{-1}$), it is very likely they will merge by $z \sim 0.7$.

5. DISCUSSION

The physical properties of the galaxies making up MS1054's bound red pairs indicate that the pairs can evolve into massive E/S0 members by $z \sim 0.7$. The color-magnitude diagram (Fig. 3) shows that 1) the pair galaxies span the range in luminosity and include the brightest cluster members; 2) they are as red (old) as the E/S0 population; and 3) their scatter in $(B - K_s)$ color is only ~ 1.5 times that of the E/S0's. Their relatively small scatter compared to, e.g. MS1054's late-types, indicates that the pair galaxies have homogeneous stellar populations; assuming these systems evolve passively, their fractional age differences will decrease and their scatter reduce accordingly. Their co-added spectrum confirms that the pair galaxies are dominated by old stellar populations (Fig. 4). The pair galaxies also include very massive members: 9 have internal velocity dispersions > 200 km s $^{-1}$ (Wuyts et al. 2004).

Can the number of E/S0 members formed via late-time merging between red members significantly alter MS1054's stellar mass function? From their study covering a wide range in environment, Bell et al. (2004) claim that the stellar mass function for red galaxies increases by a factor of two between $z \sim 1$ and now. If true, adopting the simplest scenario implies that every present-day cluster early-type undergoes a merger at $z \lesssim 1$. Assuming a typical merger timescale of 0.5 Gyr and $\Delta t_{z=1} = 7.7$ Gyr, this requires that at a given time, only $\sim 6.5\%$ of the galaxies need be in a merging system. The merger fraction in MS1054

is more than twice this value (16%), and MS1054 has a rather astonishing merger rate of ~ 20 Gyr $^{-1}$.

Although MS1054's current merger rate is certainly high, the key question is how long can it be sustained? To increase the stellar mass function of red galaxies by the quoted factor of two, MS1054 would need to retain a merger fraction of $\sim 16\%$ over the next ~ 3 Gyr (assuming $\Delta t_{MGR} = 0.5$ Gyr) and have all of these mergers evolve into early-type members. MS1054 does have a reservoir of possible "future" mergers: there are six red galaxy pairs with $30 < R_s < 50h^{-1}$ kpc and $\Delta v \leq 300$ km s $^{-1}$ (Fig. 1, right) that can provide at least one more generation of merging pairs.

Accretion of galaxy groups also can be a source of future mergers: the group environment is conducive to galaxy-galaxy merging, and Press-Schechter theory (Press & Schechter 1974) predicts MS1054 will more than double its mass by $z \sim 0$ (M. Santos, private communication) via accretion of, e.g. groups (Gonzalez et al. 2005). MS1054's high merger rate is very likely linked to its ongoing accretion of galaxy groups (Hoekstra et al. 2000); if so, this implies that an elevated merger rate is associated with this special phase in cluster evolution. MS1054's merger rate will drop as it virializes (Makino & Hut 1997), a result consistent with the low merger fractions found in lower redshift clusters (e.g. Couch et al. 1998).

Thus in the case of MS1054, it is quite feasible that all of the early-type members evolved from (passive) galaxy-galaxy mergers at $z \lesssim 1$, assuming the cluster does not virialize immediately after the epoch of observation.

The authors thank D. Fabricant for help with the visual classifications, D. Magee for help with the cluster redshift survey, and M. Postman for use of the reduced ACS imaging. K. Tran acknowledges support from the Swiss National Science Foundation; she also thanks P. Norberg and M. Santos for helpful discussions. Additional support from HST-GO09803.05-A and NAG5-7697 also are acknowledged. Finally, the authors extend special thanks to those of Hawaiian ancestry on whose sacred mountain we are privileged to be guests.

REFERENCES

- Barnes, J. E. 1988, *ApJ*, 331, 699
 Bell, E. F., Wolf, C., Meisenheimer, K., Rix, H., Borch, A., Dye, S., Kleinheinrich, M., Wisotzki, L., & McIntosh, D. H. 2004, *ApJ*, 608, 752
 Bertin, E. & Arnouts, S. 1996, *A&A*, 117, 393
 Couch, W. J., Barger, A. J., Smail, I., Ellis, R. S., & Sharples, R. M. 1998, *ApJ*, 497, 188
 Dressler, A. 1980, *ApJ*, 236, 351
 Förster Schreiber, N. M., van Dokkum, P. G., Franx, M., Labbé, I., Rudnick, G., & Daddi, E. 2005, *AJ*, submitted
 Gonzalez, A. H., Tran, K. H., Conbere, M. N., & Zaritsky, D. 2005, *ApJ*, 624, L73
 Hoekstra, H., Franx, M., & Kuijken, K. 2000, *ApJ*, 532, 88
 Jonsson, P., Cox, T., Primack, J., & Somerville, R. 2005, *ApJ*
 Kelson, D. D., Illingworth, G. D., Franx, M., & van Dokkum, P. G. 2001, *ApJ*, 552, L17
 Lin, L., Koo, D. C., Willmer, C. N. A., Patton, D. R., Conselice, C. J., Yan, R., Coil, A. L., Cooper, M. C., Davis, M., Faber, S. M., Gerke, B. F., Guhathakurta, P., & Newman, J. A. 2004, *ApJ*, 617, L9
 Makino, J. & Hut, P. 1997, *ApJ*, 481, 83
 Mihos, J. C. & Hernquist, L. 1994, *ApJ*, 431, L9
 Oke, J. B., Cohen, J. G., Carr, M., Cromer, J., Dingizian, A., Harris, F. H., Labrecque, S., Luciano, R., Schaal, W., Epps, H., & Miller, J. 1995, *PASP*, 107, 375
 Patton, D. R., Carlberg, R. G., Marzke, R. O., Pritchet, C. J., da Costa, L. N., & Pellegrini, P. S. 2000, *ApJ*, 536, 153
 Postman, M., Franx, M., Cross, N., Holden, B., & Ford, H. 2005, *ApJ*, in press
 Press, W. H. & Schechter, P. 1974, *ApJ*, 187, 425
 Springel, V. 2000, *MNRAS*, 312, 859
 Tran, K. H., Franx, M., Illingworth, G., Kelson, D. D., & van Dokkum, P. 2003, *ApJ*, 599, 865
 Tran, K. H., Kelson, D. D., van Dokkum, P., Franx, M., Illingworth, G. D., & Magee, D. 1999, *ApJ*, 522
 van Dokkum, P. G., Franx, M., Fabricant, D., Illingworth, G. D., & Kelson, D. D. 2000a, *ApJ*, 541, 95
 van Dokkum, P. G., Franx, M., Fabricant, D., Kelson, D. D., & Illingworth, G. D. 1999, *ApJ*, 520, L95
 van Dokkum, P. G., Franx, M., Kelson, D. D., Illingworth, G. D., & Fabricant, D. 2000b, *astro-ph/0012090*, 541
 Wuyts, S., van Dokkum, P. G., Kelson, D. D., Franx, M., & Illingworth, G. D. 2004, *ApJ*, 605, 677




FIG. 1.— *Left*: The relative line-of-sight velocity (Δv) of six merging galaxy pairs from vD99 (hatched) compared to the normalized distribution of 10^4 galaxy pairs randomly drawn from the full cluster sample; all six merging pairs have projected separations of $R_s < 10h^{-1}$ kpc. A K-S test confirms the two distributions differ at the $> 99.7\%$ level, thus the merging pairs are not due to chance projections between members and are bound systems. *Middle*: The relative galaxy density (δ) as a function of increasing galaxy-galaxy separation (R_s) in MS1054. Due to the high galaxy densities in the cluster core, we exclude the inner $R_{BCG} < 200h^{-1}$ kpc region. MS1054 has a strong peak at $R_s < 10h^{-1}$ kpc; this excess of pairs at small separations independently confirms that MS1054 has an unusually large number of close red pairs. *Right*: Relative velocity (Δv) as a function of projected separation (R_s) for cluster pairs where redshifts of both galaxies have been measured. We consider a bound pair to be a system composed of two galaxies with $\Delta v \leq 300$ km s $^{-1}$ and $R_s < 30h^{-1}$ kpc (solid lines): there are 10 such pairs. MS1054 also has six red galaxy pairs with $30 < R_s < 50h^{-1}$ kpc (dotted line) and $\Delta v \leq 300$ km s $^{-1}$; these are likely to be future mergers.




FIG. 2.— [Higher resolution version available at <http://www.exp-astro.phys.ethz.ch/tran/outgoing/ms1054mgrs.ps.gz>] HST/ACS images ($5'' \times 5''$) of the 10 objectively selected, spectroscopically confirmed bound red galaxy pairs ($R_s \leq 30h^{-1}$ kpc, $\Delta v \leq 300$ km s $^{-1}$). The five red pairs that were classified as mergers by vD99 are H3495/3524 (#1340), H4459/4528 (#1692), H5347/5543 (#1760), H5840/5940 (#1583), and H6036/6064 (#1163).




FIG. 3.— Color-magnitude (CM) diagram binned by galaxy type. The CM relation shown in each panel is determined by fitting an iterative least-squares to the E/S0 members (solid line). Galaxies in bound red pairs span the range in luminosity and include the brightest cluster members. We also list the deviation from the CM relation where we have measured the median offset (C_{MED}) and mean absolute deviation in color (σ_{MAD} ; “scatter”). The pair galaxies are as red as the E/S0 members and their scatter in color is relatively small, especially compared to the late-type members.




FIG. 4.— Smoothed co-added spectra of cluster members in bound red pairs; the vertical dotted lines denote $[OII]\lambda 3727$ and $H\delta$. All spectra are normalized by their flux at $4000\text{\AA}$ before being co-added. The strong break at $4000\text{\AA}$ is characteristic of a stellar population dominated by old (G & K) stars.